


MIT News

ON CAMPUS AND AROUND THE WORLD



A brain slice that shows the hippocampus cells that were activated by the new stimulation technique (bottom image, lighter green areas on the left)

Courtesy of the researchers

A noninvasive method for deep brain stimulation

Electrodes placed on the scalp could help patients with brain diseases.

Anne Trafton | MIT News Office
June 1, 2017

Delivering an electrical current to a part of the brain involved in movement control has proven successful in treating many Parkinson's disease patients. This approach, known as deep brain stimulation, requires implanting electrodes in the brain — a complex procedure that carries some risk to the patient.

Now, MIT researchers, collaborating with investigators at Beth Israel Deaconess Medical Center (BIDMC) and the IT'IS Foundation, have come up with a way to stimulate regions deep within the brain using electrodes placed on the scalp. This approach could make deep brain stimulation noninvasive, less risky, less expensive, and more accessible to patients.

"Traditional deep brain stimulation requires opening the skull and implanting an electrode, which can have complications. Secondly, only a small number of people can do this kind of neurosurgery," says Ed Boyden, an associate professor of biological engineering and brain and cognitive sciences at MIT, and the senior author of the study, which appears in the June 1 issue of *Cell*.

Doctors also use deep brain stimulation to treat some patients with obsessive compulsive disorder, epilepsy, and depression, and are exploring the possibility of using it to treat other conditions such as autism. The new, noninvasive approach could make it easier to adapt deep brain stimulation to treat additional disorders, the researchers say.

PRESS MENTIONS

MIT researchers have developed a noninvasive method to stimulate specific neurons deep in the brain that could be used to help treat patients with diseases such as Parkinson's, reports Meredith Wadman for *Science*. This new method could also allow scientists to "selectively prod deep-brain neurons into action," explains Wadman.

Science

Writing for *Wired*, Abigail Beal highlights how MIT researchers have developed a noninvasive technique to trigger reactions in deep brain cells using low frequency electrical signals. "If we could noninvasively stimulate deep regions, without hitting overlying regions, we might be able to help more people because we could stimulate deep regions selectively, without needing surgery," explains Prof. Ed Boyden.

WIRED

“With the ability to stimulate brain structures noninvasively, we hope that we may help discover new targets for treating brain disorders,” says the paper’s lead author, Nir Grossman, a former Wellcome Trust-MIT postdoc working at MIT and BIDMC, who is now a research fellow at Imperial College London.

Deep locations

Electrodes for treating Parkinson’s disease are usually placed in the subthalamic nucleus, a lens-shaped structure located below the thalamus, deep within the brain. For many Parkinson’s patients, delivering electrical impulses in this brain region can improve symptoms, but the surgery to implant the electrodes carries risks, including brain hemorrhage and infection.

Other researchers have tried to noninvasively stimulate the brain using techniques such as transcranial magnetic stimulation (TMS), which is FDA-approved for treating depression. Since TMS is noninvasive, it has also been used in normal human subjects to study the basic science of cognition, emotion, sensation, and movement. However, using TMS to stimulate deep brain structures can also result in surface regions being strongly stimulated, resulting in modulation of multiple brain networks.

The MIT team devised a way to deliver electrical stimulation deep within the brain, via electrodes placed on the scalp, by taking advantage of a phenomenon known as temporal interference.

This strategy requires generating two high-frequency electrical currents using electrodes placed outside the brain. These fields are too fast to drive neurons. However, these currents interfere with one another in such a way that where they intersect, deep in the brain, a small region of low-frequency current is generated inside neurons. This low-frequency current can be used to drive neurons’ electrical activity, while the high-frequency current passes through surrounding tissue with no effect.

By tuning the frequency of these currents and changing the number and location of the electrodes, the researchers can control the size and location of the brain tissue that receives the low-frequency stimulation. They can target locations deep within the brain without affecting any of the surrounding brain structures. They can also steer the location of stimulation, without moving the electrodes, by altering the currents. In this way, deep targets could be stimulated, both for therapeutic use and basic science investigations.

“You can go for deep targets and spare the overlying neurons, although the spatial resolution is not yet as good as that of deep brain stimulation,” says Boyden, who is a member of MIT’s Media Lab and McGovern Institute for Brain Research.

“The fact that only two sources are needed and one can conceptually steer the focus is fascinating. I can see a possible future where this technology could be made portable for non-invasive deep brain stimulation as a therapy. If the work could be shown to do so in larger animal models than a mouse I would be even more optimistic,” says Tipu Aziz, a professor of neurosurgery at Oxford University, who was not involved in the research.

Targeted stimulation

NBC News reporter Maggie Fox writes that MIT researchers have developed a noninvasive brain stimulation technique that could eventually offer relief to patients with diseases like Parkinson’s and epilepsy without requiring surgery. Fox explains that the method allows for sending, “electrical signals deep into the brain without affecting the layers in between.”



New York Times reporter Pam Belluck writes that MIT researchers have developed a new, non-invasive deep brain stimulation technique. The technique could be used to help treat, “a range of neurological and psychiatric disorders more cheaply and safely than current approaches,” writes Belluck.



MIT researchers have developed a non-invasive technique for deep brain stimulation, which could be used to help patients with brain diseases, reports Mo Costandi for *The Guardian*. “Targets for disorders such as depression, Alzheimer’s, PTSD, and so forth, are deep in the brain, and they might be more selectively stimlatable with our method,” says Prof. Ed Boyden.



RELATED

- Ed Boyden
- Li-Huei Tsai
- Department of Brain and Cognitive Sciences
- Department of Biological Engineering
- McGovern Institute for Brain Research
- Picower Institute for Learning and Memory
- MIT Media Lab
- School of Engineering

Li-Huei Tsai, director of MIT's Picower Institute for Learning and Memory, and researchers in her lab tested this technique in mice and found that they could stimulate small regions deep within the brain, including the hippocampus. They were also able to shift the site of stimulation, allowing them to activate different parts of the motor cortex and prompt the mice to move their limbs, ears, or whiskers.

"We showed that we can very precisely target a brain region to elicit not just neuronal activation but behavioral responses," says Tsai, who is an author of the paper. "I think it's very exciting because Parkinson's disease and other movement disorders seem to originate from a very particular region of the brain, and if you can target that, you have the potential to reverse it."

Significantly, in the hippocampus experiments, the technique did not activate the neurons in the cortex, the region lying between the electrodes on the skull and the target deep inside the brain. The researchers also found no harmful effects in any part of the brain.

Last year, Tsai showed that using light to visually induce brain waves of a particular frequency could substantially reduce the beta amyloid plaques seen in Alzheimer's disease, in the brains of mice. She now plans to explore whether this type of electrical stimulation could offer a new way to generate the same type of beneficial brain waves.

Other authors of the paper are MIT research scientist David Bono; former MIT postdocs Suhasa Kodandaramaiah and Andrii Rudenko; MIT postdoc Nina Dedic; MIT grad student Ho-Jun Suk; Beth Israel Deaconess Medical Center and Harvard Medical School Professor Alvaro Pascual-Leone; and ETH Zurich researchers Antonino Cassara, Esra Neufeld, and Niels Kuster.

The research was funded in part by the Wellcome Trust, a National Institutes of Health Director's Pioneer Award, an NIH Director's Transformative Research Award, the New York Stem Cell Foundation Robertson Investigator Award, the MIT Center for Brains, Minds, and Machines, Jeremy and Joyce Wertheimer, Google, a National Science Foundation Career Award, the MIT Synthetic Intelligence Project, and Harvard Catalyst: The Harvard Clinical and Translational Science Center.

School of Science

School of Architecture and Planning


ARCHIVES



Unique visual stimulation may be new treatment for Alzheimer's



Noninvasive brain control



Illuminating neuron activity in 3-D

-
- Topics: [Research](#) [Parkinson's](#) [Brain and cognitive sciences](#) [Media Lab](#)
[Biological engineering](#) [McGovern Institute](#) [Picower Institute](#)
[School of Science](#) [School of Engineering](#)
[School of Architecture and Planning](#)

About This Website

This Website is maintained by the MIT News Office, part of the Office of Communications.

MIT News Office • Building 11-400
Massachusetts Institute of Technology • Cambridge, MA 02139-4307